

A. Diabetes




Figure 1A-H. Cost ratio estimates (with 95% confidence intervals) for predictors of total health care costs for 8 chronic conditions in fiscal year 2010-2011.^a

^a Cost ratios (Exp[B]) were estimated using generalized linear models with the log link function and γ -distributed errors and all predictors entered simultaneously. Error bars indicate 95% confidence intervals.

FTE = full-time equivalent; RUB = Resource Utilization Band.

B. CHF




Figure 1A-H. Cost ratio estimates (with 95% confidence intervals) for predictors of total health care costs for 8 chronic conditions in fiscal year 2010-2011.^a

^a Cost ratios (Exp[B]) were estimated using generalized linear models with the log link function and γ -distributed errors and all predictors entered simultaneously. Error bars indicate 95% confidence intervals.

CHF = congestive heart failure; FTE = full-time equivalent; RUB = resource utilization band.

C. COPD




Figure 1A-H. Cost ratio estimates (with 95% confidence intervals) for predictors of total health care costs for 8 chronic conditions in fiscal year 2010-2011.^a

^a Cost ratios (Exp[B]) were estimated using generalized linear models with the log link function and γ -distributed errors and all predictors entered simultaneously. Error bars indicate 95% confidence intervals.

COPD = chronic obstructive pulmonary disease; FTE = full-time equivalent; RUB = resource utilization band.

D. Hypertension




Figure 1A-H. Cost ratio estimates (with 95% confidence intervals) for predictors of total health care costs for 8 chronic conditions in fiscal year 2010-2011.^a

^a Cost ratios (Exp[B]) were estimated using generalized linear models with the log link function and γ -distributed errors and all predictors entered simultaneously. Error bars indicate 95% confidence intervals.

FTE = full-time equivalent; RUB = resource utilization band.

E. Angina




Figure 1A-H. Cost ratio estimates (with 95% confidence intervals) for predictors of total health care costs for 8 chronic conditions in fiscal year 2010-2011.^a

^a Cost ratios [Exp(B)] were estimated using generalized linear models with the log link function and γ -distributed errors and all predictors entered simultaneously. Error bars indicate 95% confidence intervals.

FTE = full-time equivalent; RUB = Resource Utilization Band.

F. CKD




Figure 1A-H. Cost ratio estimates (with 95% confidence intervals) for predictors of total health care costs for 8 chronic conditions in fiscal year 2010-2011.^a

^a Cost ratios (Exp[B]) were estimated using generalized linear models with the log link function and γ -distributed errors and all predictors entered simultaneously. Error bars indicate 95% confidence intervals.

CKD = chronic kidney disease; FTE = full-time equivalent; RUB = Resource Utilization Band.

G. Osteoarthritis




Figure 1A-H. Cost ratio estimates (with 95% confidence intervals) for predictors of total health care costs for 8 chronic conditions in fiscal year 2010-2011.^a

^a Cost ratios (Exp[B]) were estimated using generalized linear models with the log link function and γ -distributed errors and all predictors entered simultaneously. Error bars indicate 95% confidence intervals.

FTE = full-time equivalent; RUB = Resource Utilization Band.

H. Stroke




Figure 1A-H. Cost ratio estimates (with 95% confidence intervals) for predictors of total health care costs for 8 chronic conditions in fiscal year 2010-2011.^a

^a Cost ratios (Exp[B]) were estimated using generalized linear models with the log link function and γ -distributed errors and all predictors entered simultaneously. Error bars indicate 95% confidence intervals.

FTE = full-time equivalent; RUB = Resource Utilization Band.